

ON THE EQUIVARIANT COHOMOLOGY OF SUBVARIETIES OF A $\mathfrak{B}$ -REGULAR VARIETY

JAMES B. CARRELL AND KIUMARS KAVEH

To Bert Kostant on his 80th birthday.

Abstract. By a $\mathfrak{B}$ -regular variety, we mean a smooth projective variety over $\mathbb{C}$ admitting an algebraic action of the upper triangular Borel subgroup $\mathfrak{B} \subset SL_2(\mathbb{C})$ such that the unipotent radical in $\mathfrak{B}$ has a unique fixed point. A result of M. Brion and the first author [4] describes the equivariant cohomology algebra (over $\mathbb{C}$) of a $\mathfrak{B}$ -regular variety X as the coordinate ring of a remarkable affine curve in $X \times \mathbb{P}^1$. The main result of this paper uses this fact to classify the $\mathfrak{B}$ -invariant subvarieties Y of a $\mathfrak{B}$ -regular variety X for which the restriction map $i_Y^* : H^*(X) \rightarrow H^*(Y)$ is surjective.

1. INTRODUCTION

A nilpotent element e in the Lie algebra $\mathfrak{g}$ of a complex semi-simple Lie group G is *regular* if it lies in a unique Borel subalgebra $\mathfrak{b}$ of $\mathfrak{g}$. If we let B be the unique Borel subgroup of G with $\text{Lie}(B) = \mathfrak{b}$ and recall that the flag variety G/B of G parameterizes the family of all Borel subalgebras of $\mathfrak{g}$, it follows that the one parameter group $\exp(te)$ ($t \in \mathbb{C}$) of G acts on G/B by left translation with unique fixed point the identity coset B or, equivalently, the unique Borel subalgebra $\mathfrak{b}$ containing e . By the Jacobson-Morosov Lemma, e determines a two dimensional solvable subalgebra $\mathfrak{e}$ of $\mathfrak{g}$ isomorphic to the Lie algebra of the upper triangular Borel subgroup $\mathfrak{B}$ of $SL_2(\mathbb{C})$. The two dimensional solvable subgroup E of G determined by $\mathfrak{e}$ is thus isomorphic to a Borel either in $SL_2(\mathbb{C})$ or $PSL_2(\mathbb{C})$ and hence is a homomorphic image of $\mathfrak{B}$. Consequently, $\mathfrak{B}$ acts on G/B , via E , such that its unipotent radical $\mathfrak{U}$ has exactly one fixed point, namely B .

In view of this, one may generalize the notion of a regular nilpotent by considering an algebraic action $\mathfrak{B} \curvearrowright X$ of the upper triangular group $\mathfrak{B}$ on a smooth complex projective variety X such that its unipotent radical $\mathfrak{U}$ has a unique fixed point $o \in X$. For such an action, the maximal torus $\mathfrak{T}$ on the diagonal of $\mathfrak{B}$ is known to have a finite fixed point set (see [5]). In [4, 5], such an action is called *regular* and X is called a *regular variety*. Slightly changing this terminology, we will henceforth call the action $\mathfrak{B}$ -regular and say that X is a *$\mathfrak{B}$ -regular variety*. By the above remarks, the flag variety G/B and, more generally, all algebraic homogeneous spaces G/P , P a parabolic in G , give a rich class of $\mathfrak{B}$ -regular varieties.

The main goal of this paper is to study the cohomology algebras of $\mathfrak{B}$ -invariant subvarieties of a $\mathfrak{B}$ -regular variety. For example, Schubert varieties in a G/B or G/P (that is, closures of B -orbits) form an important class of

examples of $\mathfrak{B}$ -invariant subvarieties to which our results will apply. Moreover, certain nilpotent Hessenberg varieties (including the Peterson variety) [12] and certain Springer fibres corresponding to nilpotents in the centralizer of a regular nilpotent $e \in \mathfrak{g}$ give further interesting examples of $\mathfrak{B}$ -invariant subvarieties in G/B which we hope to investigate in a future work.

If X is a $\mathfrak{B}$ -regular variety, then one knows the remarkable fact that its cohomology algebra $H^*(X)$ over $\mathbb{C}$ is isomorphic with the coordinate ring $A(Z)$ of the zero scheme Z associated to the algebraic vector field on X generated by $\mathfrak{U}$ (cf. [1, 2, 5]). Moreover, its $\mathfrak{T}$ -equivariant cohomology $H_{\mathfrak{T}}^*(X)$ is isomorphic to the coordinate ring of a canonical $\mathfrak{T}$ -stable affine curve $\mathcal{Z}_X$ in $X \times \mathbb{P}^1$ ([4]). That is, $\text{Spec}(H_{\mathfrak{T}}^*(X))$ is $\mathcal{Z}_X$. These isomorphisms extend to what we will call *principal subvarieties*: namely, $\mathfrak{B}$ -invariant subvarieties Y of X for which the natural restriction map $i_Y^* : H^*(X) \rightarrow H^*(Y)$ is surjective (cf. Theorem 1). In particular, if Y is principal, then $\text{Spec}(H_{\mathfrak{T}}^*(Y))$ is the union $\mathcal{Z}_Y$ of the irreducible components of $\mathcal{Z}_X$ lying over $Y \times \mathbb{C}$. Furthermore, if $Y \cap Z$ denotes the schematic intersection of Y and Z , then the coordinate ring $A(Y \cap Z)$ is isomorphic to $H^*(Y)$ as long as $\dim A(Y \cap Z) = \dim H^*(Y)$. With these facts as motivation, our aim is to classify the principal subvarieties of a $\mathfrak{B}$ -regular variety. This will follow from a general result which describes the image of $H_{\mathfrak{T}}^*(X)$ in $H_{\mathfrak{T}}^*(Y)$ under i_Y^* for any $\mathfrak{B}$ -invariant subvariety Y of a $\mathfrak{B}$ -regular variety X .

The curve $\mathcal{Z}_X$ admits a natural description. Consider the diagonal action of $\mathfrak{B}$ on $X \times \mathbb{P}^1$, where $\mathfrak{B}$ acts on $\mathbb{P}^1$ by the standard action of $SL_2(\mathbb{C})$ on $\mathbb{C}^2$. Then the irreducible components of $\mathcal{Z}_X$ have the form $\mathfrak{B} \cdot (\zeta, \infty) \setminus (\zeta, \infty)$, where ζ ranges over all of $X^{\mathfrak{T}}$ and ∞ denotes the point $[0, 1] \in \mathbb{P}^1$. The complete description of the equivariant cohomology of a $\mathfrak{B}$ -regular variety X and principal subvariety Y proved in [4] is given by

Theorem 1. (cf. [4]) *If X is a $\mathfrak{B}$ -regular variety, then there exists a graded $\mathbb{C}$ -algebra isomorphism $\rho_X : H_{\mathfrak{T}}^*(X) \rightarrow \mathbb{C}[\mathcal{Z}_X]$. Furthermore, if Y is a principal subvariety and $\mathcal{Z}_Y$ is the (reduced) affine curve $\mathcal{Z}_X \cap (Y \times \mathbb{C})$, then there is also a graded $\mathbb{C}$ -algebra isomorphism $\rho_Y : H_{\mathfrak{T}}^*(Y) \rightarrow \mathbb{C}[\mathcal{Z}_Y]$ making the diagram*

$$(1) \quad \begin{array}{ccc} H_{\mathfrak{T}}^*(X) & \xrightarrow{\rho_X} & \mathbb{C}[\mathcal{Z}_X] \\ i_Y^* \downarrow & & \downarrow \overline{i_Y^*} \\ H_{\mathfrak{T}}^*(Y) & \xrightarrow{\rho_Y} & \mathbb{C}[\mathcal{Z}_Y], \end{array}$$

commutative, where the vertical maps are natural restrictions. Moreover, the horizontal maps are $\mathbb{C}[v]$ -module maps under the standard $\mathbb{C}[v]$ -module structure on $H_{\mathfrak{T}}^(X)$ and $H_{\mathfrak{T}}^*(Y)$ and the $\mathbb{C}[v]$ -module structure on $\mathbb{C}[\mathcal{Z}_X]$ and $\mathbb{C}[\mathcal{Z}_Y]$ induced by the second projection.*

One easily sees from the definitions that if v is an affine coordinate on $\mathbb{C} = \mathbb{P}^1 \setminus [1, 0]$, then $\mathbb{C}[\mathcal{Z}_X]/(v)\mathbb{C}[\mathcal{Z}_X] \cong A(Z)$. Hence, Theorem 1 allows one to see the isomorphism $H^*(X) \cong A(Z)$ in a natural way from elementary properties of equivariant cohomology. Indeed, it turns out that ρ_X maps the augmentation ideal $(v)H_{\mathfrak{T}}^*(X)$ in $H_{\mathfrak{T}}^*(X)$ to $(v)\mathbb{C}[\mathcal{Z}_X]$, so $A(Z) \cong H^*(X)$

since $H_{\mathfrak{T}}^*(X)/(v)H_{\mathfrak{T}}^*(X) \cong H^*(X)$. Similarly, $H^*(Y) \cong \mathbb{C}[\mathcal{Z}_Y]/(v)\mathbb{C}[\mathcal{Z}_Y]$ if Y is principal. However, it is not in general true that $\mathbb{C}[\mathcal{Z}_Y]/(v)\mathbb{C}[\mathcal{Z}_Y]$ is isomorphic to $A(Y \cap Z)$.

Since surjectivity of $i_Y^* : H^*(X) \rightarrow H^*(Y)$ is equivalent to surjectivity of $i_Y^* : H_{\mathfrak{T}}^*(X) \rightarrow H_{\mathfrak{T}}^*(Y)$, (1) suggests an approach to the surjectivity question. Namely, since $\overline{i_Y^*} : \mathbb{C}[\mathcal{Z}_X] \rightarrow \mathbb{C}[\mathcal{Z}_Y]$ is surjective for any $\mathfrak{B}$ -invariant subvariety Y , the question of surjectivity of i_Y^* boils down to determining if there exists an injective map $\rho_Y : H_{\mathfrak{T}}^*(Y) \rightarrow \mathbb{C}[\mathcal{Z}_Y]$ such that (1) is commutative. We will resolve this question by showing

Theorem 2. *A $\mathfrak{B}$ -invariant subvariety Y of a $\mathfrak{B}$ -regular variety X is principal if and only if $H_{\mathfrak{T}}^*(Y)$ (equivalently, $H^*(Y)$) is generated by Chern classes of $\mathfrak{B}$ -equivariant algebraic vector bundles on Y .*

An example of a $\mathfrak{B}$ -invariant subvariety which is not principal is easily obtained. In fact, let X be $\mathfrak{B}$ -regular, and let Y denote the union of all the $\mathfrak{B}$ -stable curves in X . Then $X^{\mathfrak{T}} = Y^{\mathfrak{T}}$, and it is not hard to see that i_Y^* is not surjective if $\dim X > 1$. However, $H_{\mathfrak{T}}^*(Y)$ is not generated by $\mathfrak{B}$ -equivariant vector bundles. We will verify this claim for $X = \mathbb{P}^2$ in Section 5.

Schubert varieties in G/B or a G/P are well known to be principal. Indeed, the B -orbits are affine cells, and every Schubert variety is a union of B -orbits. Moreover, certain Springer fibres in $SL_n(\mathbb{C})/B$ are principal. To see this, first recall that a Springer fibre in G/B is by definition the fixed point set of a unipotent element of G . Viewing G/B as the variety of Borel subalgebras of $\mathfrak{g}$, this definition is equivalent defining a Springer fibre to be the set of all Borels in $\mathfrak{g}$ containing a fixed nilpotent in $\mathfrak{g}$. By classical result of Spaltenstein [11], the cohomology map $H^*(G/B) \rightarrow H^*(Y)$ is surjective for every Springer fibre Y provided $G = SL_n(\mathbb{C})$. Let e denote a regular nilpotent in $\text{Lie}(B)$, and let $\mathfrak{B} \circ G/B$ denote the regular action determined by e as explained in the first paragraph. Any Springer fibre in G/B corresponding to a nilpotent in the centralizer $\mathfrak{g}^e$ of e which is also a $\mathfrak{T}$ -weight vector is $\mathfrak{B}$ -invariant. Thus such Springer fibres form a class of principal subvarieties of $SL_n(\mathbb{C})/B$. This example shows that principal subvarieties need not be irreducible.

2. PRELIMINARIES

Let $\lambda : \mathbb{C}^* \rightarrow \mathfrak{T}$ and $\varphi : \mathbb{C} \rightarrow \mathfrak{U}$ denote the one parameter subgroups

$$\lambda(t) = \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix}, \quad \varphi(v) = \begin{pmatrix} 1 & v \\ 0 & 1 \end{pmatrix}.$$

Then $\lambda(t)\varphi(v)\lambda(t)^{-1} = \varphi(t^2v)$ for all $t \in \mathbb{C}^*$ and $v \in \mathbb{C}$. Suppose X is a $\mathfrak{B}$ -regular variety with $X^{\mathfrak{U}} = \{o\}$, and note that $o \in X^{\mathfrak{T}}$. Put

$$X_o = \{x \in X \mid \lim_{t \rightarrow \infty} \lambda(t) \cdot x = o\}.$$

Recall from [2] (also see [5]) that X_o is an open neighborhood of o isomorphic with $T_o(X)$. Thus, if $a_1, \dots, a_n$ ($n = \dim X$) are the weights of λ on

$T_o(X)$ repeated with multiplicities, then all $a_i < 0$ and there exist affine coordinates $u_1, \dots, u_n$ on X_o which are quasi-homogeneous of positive degrees $-a_1, \dots, -a_n$ with respect to λ . That is,

$$\lambda(t) \cdot u_i = t^{-a_i} u_i,$$

for all i . The induced positive grading on $\mathbb{C}[X_o] = \mathbb{C}[u_1, \dots, u_n]$ is frequently called the *principal grading*. Since the fixed point set $X^{\mathfrak{T}}$ of the torus $\mathfrak{T}$ is finite and contains o ([2]), we will write $X^{\mathfrak{T}} = \{\zeta_1, \dots, \zeta_r\}$, where $\zeta_1 = o$.

Note that the natural action of $SL_2(\mathbb{C})$ on $\mathbb{P}^1$ induces a regular action. If $\mathfrak{B}$ denotes the upper triangular matrices and $\mathfrak{T}$ the diagonals, then $\zeta_1 = [1, 0]$, $\zeta_2 = [0, 1]$ and the big cell is $\{[1, v] \mid v \in \mathbb{C}\}$.

From now on, $\mathfrak{B}$ will denote the upper triangular Borel in $SL_2(\mathbb{C})$ and $\mathfrak{T}$ will be the diagonal torus; X will always denote a $\mathfrak{B}$ -regular variety. To simplify the notation, we will put $0 = [1, 0]$ and $\infty = [0, 1]$ so $\mathbb{P}^1 = \mathbb{C} \cup \{\infty\}$. Notice that the diagonal action $\mathfrak{B} \curvearrowright (X \times \mathbb{P}^1)$ is also regular with $\mathfrak{U}$ -fixed point $(o, 0)$. Define a projective curve $Z_X \subset X \times \mathbb{P}^1$ as follows: let Z_i be Zariski closure of the orbit $\mathfrak{B}(\zeta_i, \infty)$, where $i \geq 1$, and let

$$Z_X = \bigcup_{1 \leq i \leq r} Z_i.$$

Thus Z_X is a $\mathfrak{B}$ -stable curve with $r = |X^{\mathfrak{T}}|$ irreducible components. Moreover, the second projection $p_2 : X \times \mathbb{P}^1 \rightarrow \mathbb{P}^1$ induces an isomorphism on each component. Hence Z_X is a bouquet of r $\mathbb{P}^1$ s passing through $(o, 0)$. Now define an affine curve

$$(2) \quad \mathcal{Z}_X = Z_X \cap (X_o \times \mathbb{C}),$$

where the intersection is assumed to be in the sense of varieties, i.e. reduced. This curve is $\mathfrak{T}$ -stable but not $\mathfrak{B}$ -stable. In fact, $\mathcal{Z}_X$ is obtained from Z_X by removing the point at infinity on each irreducible component. In particular, the coordinate ring $\mathbb{C}[\mathcal{Z}_X]$ is a graded $\mathbb{C}$ -algebra via the principal grading, and the projection p_2 induces a $\mathbb{C}[v]$ -module structure on $\mathbb{C}[\mathcal{Z}_X]$, where v denotes the affine coordinate on $\mathbb{C}$. For later use, we observe (cf. [4, p.192])

$$(3) \quad \text{if } v \neq 0, \text{ then } (x, v) \in \mathcal{Z}_X \text{ if and only if } \varphi(-v^{-1}) \cdot x \in X^{\mathfrak{T}}.$$

We now recall the basic isomorphism $\rho_X : H_{\mathfrak{T}}^*(X) \rightarrow \mathbb{C}[\mathcal{Z}_X]$. Since the odd cohomology of a $\mathfrak{B}$ -regular variety is trivial, the localization theorem in equivariant cohomology implies $i^* : H_{\mathfrak{T}}^*(X) \rightarrow H_{\mathfrak{T}}^*(X^{\mathfrak{T}})$ is injective, where $i : X^{\mathfrak{T}} \rightarrow X$ is the inclusion map. Thus, to each $\alpha \in H_{\mathfrak{T}}^*(X)$, one may assign an r -tuple $(\alpha_1, \dots, \alpha_r) \in \bigoplus_i \mathbb{C}[v]$. Now define a function $\rho_X(\alpha)$ on $\mathcal{Z}_X$ by putting

$$(4) \quad \rho_X(\alpha)(x, v) = \alpha_i(v),$$

if $(x, v) \in Z_i$. A key fact is that $\rho_X(\alpha)$ is a regular function on $\mathcal{Z}_X$.

3. SOME APPLICATIONS OF SURJECTIVITY

We will now use Theorem 1 to draw some conclusions about surjectivity. Suppose that Y is another $\mathfrak{B}$ -regular variety and $F : Y \rightarrow X$ a $\mathfrak{B}$ -equivariant map. We claim F induces a $\mathfrak{T}$ -equivariant map $\overline{F} : \mathcal{Z}_Y \rightarrow \mathcal{Z}_X$ by putting

$\overline{F}(y, v) = (F(y), v)$. For $F(Y^{\mathfrak{T}}) \subset X^{\mathfrak{T}}$, so if $(y, v) \in \mathcal{Z}_Y$, say $(y, v) = (b \cdot y_i, b \cdot \infty)$ for some $y_i \in Y^{\mathfrak{T}}$ and $b \in \mathfrak{B}$, we see that

$$(F(y), v) = (F(b \cdot y_i), b \cdot \infty) = (b \cdot F(y_i), b \cdot \infty) = b \cdot (F(y_i), \infty) \in \mathcal{Z}_X.$$

But $b \cdot (F(y_i), \infty) \neq (F(y_i), \infty)$ unless $b = 1$, hence the claim. Thus there is a commutative diagram

$$(5) \quad \begin{array}{ccc} H_{\mathfrak{T}}^*(X) & \xrightarrow{\rho_X} & \mathbb{C}[\mathcal{Z}_X] \\ F^* \downarrow & & \overline{F}^* \downarrow \\ H_{\mathfrak{T}}^*(Y) & \xrightarrow{\rho_Y} & \mathbb{C}[\mathcal{Z}_Y]. \end{array}$$

We now derive some consequences of this.

Theorem 3. *Assume $F : Y \rightarrow X$ is a $\mathfrak{B}$ -equivariant map of $\mathfrak{B}$ -regular varieties such that the differential dF_o of F at o is injective. Suppose also that $F|Y^{\mathfrak{T}}$ is injective. Then the restriction map $F^* : H_{\mathfrak{T}}^*(X) \rightarrow H_{\mathfrak{T}}^*(Y)$ is surjective. Consequently, $F^* : H^*(X) \rightarrow H^*(Y)$ is also surjective.*

Proof. First suppose Y is a $\mathfrak{B}$ -invariant subvariety of X which is also $\mathfrak{B}$ -regular and $F = i_Y$. By Theorem 1, the morphism ρ_Y in the diagram (5) is an isomorphism. Thus i_Y^* is surjective. Now consider the general case. Since dF_o is injective, it follows that $F(Y)$ is smooth at o . By the Borel Fixed Point Theorem applied to the singular locus of $F(Y)$, it follows that $F(Y)$ is smooth, hence $\mathfrak{B}$ -regular. Hence $i_{F(Y)}^* : H_{\mathfrak{T}}^*(X) \rightarrow H_{\mathfrak{T}}^*(F(Y))$ is surjective. As F is injective on $Y^{\mathfrak{T}}$ and $\mathfrak{B}$ -equivariant, it follows that $\overline{F} : \mathcal{Z}_Y \rightarrow \mathcal{Z}_{F(Y)}$ is also injective. Since F has a local holomorphic inverse G in a neighborhood of o , G in fact induces (by equivariant extension) a holomorphic map $\overline{G} : \mathcal{Z}_{F(Y)} \rightarrow \mathcal{Z}_Y$ which is an inverse (in the analytic category) to $\overline{F}$. Note that $F(Y)^{\mathfrak{T}} = F(Y^{\mathfrak{T}})$; that is, $F|Y^{\mathfrak{T}}$ is a bijection with $F(Y)^{\mathfrak{T}}$. For, if $w \in F(Y)^{\mathfrak{T}}$, then, by the Borel Fixed Point Theorem, the subvariety $F^{-1}(w)$ of Y contains a $\mathfrak{T}$ -fixed point due to the fact that it is $\mathfrak{T}$ -invariant. Therefore $\overline{F}^* : \mathbb{C}[\mathcal{Z}_{F(Y)}] \rightarrow \mathbb{C}[\mathcal{Z}_Y]$ is an isomorphism, giving the result. $\square$

Moreover, we also get

Theorem 4. *Two principal subvarieties of a $\mathfrak{B}$ -regular variety with the same fixed point set have isomorphic cohomology algebras (both equivariant and classical). In particular, two such subvarieties have the same dimension.*

Proof. This is an immediate consequence of Theorem 1. $\square$

Consequently, regular actions have a rather special property.

Corollary 5. *If X is $\mathfrak{B}$ -regular and Y is a $\mathfrak{B}$ -invariant subvariety of X such that $Y^{\mathfrak{T}} = X^{\mathfrak{T}}$, then either $Y = X$ or Y is singular.*

Proof. By the previous theorem, if Y is smooth, hence $\mathfrak{B}$ -regular, then $\dim Y = \dim X$. Since a $\mathfrak{B}$ -regular variety is necessarily irreducible, $Y = X$. $\square$

Of course, there are examples of torus actions on smooth projective varieties $Y \subsetneq X$ for which $X^T = Y^T$ and the conclusion of Corollary 5 fails. For example, if X is the wonderful compactification of a semi-simple algebraic group G (over $\mathbb{C}$) with maximal torus T , then all the fixed points of the (torus) action of $T \times T$ on X lie on the unique closed $G \times G$ -orbit Y (cf. [8]). This gives a proper smooth $T \times T$ -stable subvariety Y of X for which $X^{T \times T} = Y^{T \times T}$. It also shows that a wonderful compactification doesn't admit a regular action.

Finally, we have

Theorem 6. *Let X and Y be $\mathfrak{B}$ -regular varieties or principal subvarieties, and let $F : Y \rightarrow X$ be a $\mathfrak{B}$ -equivariant map such that $F(Y^\mathfrak{T}) = X^\mathfrak{T}$. Then $F^* : H_\mathfrak{T}^*(X) \rightarrow H_\mathfrak{T}^*(Y)$ is injective. In particular, if F is surjective, then $F(Y^\mathfrak{T}) = X^\mathfrak{T}$, so F^* is injective on equivariant cohomology.*

Proof. Since $F(Y^\mathfrak{T}) = X^\mathfrak{T}$, $\overline{F} : \mathcal{Z}_Y \rightarrow \mathcal{Z}_X$ is also surjective. Since $\mathcal{Z}_X$ and $\mathcal{Z}_Y$ are affine, it follows that the comorphism $\overline{F}^* : \mathbb{C}[\mathcal{Z}_X] \rightarrow \mathbb{C}[\mathcal{Z}_Y]$ is injective. Therefore, $F^* : H_\mathfrak{T}^*(X) \rightarrow H_\mathfrak{T}^*(Y)$ is injective. If F is surjective, then the Borel Fixed Point Theorem implies $F(Y^\mathfrak{T}) = X^\mathfrak{T}$. $\square$

Remark. If X and Y are smooth projective varieties and $F : Y \rightarrow X$ is surjective, it is well known that F^* is always injective on ordinary cohomology.

4. A REMARK ON A FORMULA OF KOSTANT AND MACDONALD

Let G be a complex semi-simple algebraic group, B a Borel subgroup of G and put $\mathfrak{b} = \text{Lie}(B)$. We will now describe an interesting class of $\mathfrak{B}$ -regular subvarieties of G/B which includes all smooth Schubert varieties. Let $e \in \mathfrak{b}$ be a regular nilpotent in $\mathfrak{g}$, and recall from Section 1 that e determines a regular action $\mathfrak{B} \curvearrowright G/B$ on the flag variety of G by left translation such that the identity coset $B \in G/B$ is the unique $\mathfrak{U}$ -fixed point. To simplify notation, let us identify $\mathfrak{B}$ with its homomorphic image in G such that $e \in \text{Lie}(\mathfrak{B})$ (see the comment in the first paragraph of the Introduction). Let $\mathfrak{h}$ denote a $\text{Lie}(\mathfrak{B})$ -submodule of $\mathfrak{g}$ containing $\mathfrak{b}$, and put $Y_\mathfrak{h} = \overline{\exp(\mathfrak{h})B}$.

Lemma 7. *$Y_\mathfrak{h}$ is a $\mathfrak{B}$ -invariant subvariety of G/B .*

Proof. Since the exponential map $\exp : \mathfrak{g} \rightarrow G$ is G -equivariant for the adjoint action of G on $\mathfrak{g}$, for any $x \in \text{Lie}(\mathfrak{B})$, we have

$$\exp(\text{Ad}(\exp(tx))y)B = \exp(tx)\exp(y)\exp(-tx)B = \exp(tx)\exp(y)B.$$

But for any $x, y \in \mathfrak{g}$, $\text{Ad}(\exp(tx))(y) = e^{\text{ad}(tx)}(y)$, so the term on the left side of the above identity is in $\exp(\mathfrak{h})B$ if $y \in \mathfrak{h}$. Thus $Y_\mathfrak{h}$ is stable under $\mathfrak{B}$. $\square$

Remark: If $\mathfrak{h}$ is B -stable, then $Y_\mathfrak{h}$ is also B -stable and hence is a Schubert variety.

When the variety $Y_\mathfrak{h}$ is smooth, it is a $\mathfrak{B}$ -regular variety and one can now apply the formula in [2] for the Poincaré polynomial of a $\mathfrak{B}$ -regular variety X to find an interesting class of polynomials associated to any complex

semisimple Lie algebra $\mathfrak{g}$. Let us first recall the formula. Let $a_1, \dots, a_n$ ($n = \dim X$) denote the weights of λ on $T_o(X)$ introduced in Section 2, and recall the a_i are negative integers. Then

$$(6) \quad P(X, t^{1/2}) = \prod_{1 \leq i \leq n} \frac{(1 - t^{-a_i+1})}{(1 - t^{-a_i})}.$$

In the case $X = Y_{\mathfrak{g}} = G/B$, the weights a_i are the negatives of the heights of the positive roots with respect to any maximal torus T of G contained in B , so we recover a well known formula

$$P(G/B, t^{1/2}) = \prod_{\alpha > 0} \frac{(1 - t^{ht(\alpha)+1})}{(1 - t^{ht(\alpha)})}$$

of Kostant and Macdonald.

5. A LEMMA AND AN EXAMPLE

We will now prove some facts about $\mathfrak{B}$ -invariant subvarieties of a $\mathfrak{B}$ -regular variety X . First note

Lemma 8. *Let Y be a $\mathfrak{B}$ -invariant subvariety of X with vanishing odd cohomology, and let $I(\mathcal{Z}_Y) \subset \mathbb{C}[\mathcal{Z}_X]$ denote the ideal of $\mathcal{Z}_Y$. Then:*

- (i) $\rho_X(H_{\mathfrak{T}}^{ev}(X, Y)) = I(\mathcal{Z}_Y)$; and
- (ii) *there exists a $\mathbb{C}[v]$ -algebra isomorphism*

$$\psi_Y : \mathbb{C}[\mathcal{Z}_Y] \rightarrow i_Y^*(H_{\mathfrak{T}}^*(X)) \subset H_{\mathfrak{T}}^*(Y).$$

In fact, $\psi_Y = i_Y^ \rho_X^{-1} (\overline{i_Y^*})^{-1}$.*

Proof. We first show that putting $\psi_Y = i_Y^* \rho_X^{-1} (\overline{i_Y^*})^{-1}$ gives a well defined map. Since $\mathcal{Z}_Y$ is the union of the irreducible components of $\mathcal{Z}_X$ which meet $Y \times \mathbb{C}$, the indeterminacy introduced by $(\overline{i_Y^*})^{-1}$ is supported on the complement of $\mathcal{Z}_Y$. Thus ψ_Y is indeed well defined since $\overline{i_Y^*}$ is surjective, and two classes in $H_{\mathfrak{T}}^*(X)$ which have the same image under $\overline{i_Y^*} \rho_X$ have the same image under i_Y^* , by the localization theorem and the definition of ρ_X .

We now show $\rho_X(H_{\mathfrak{T}}^{ev}(X, Y)) = I(\mathcal{Z}_Y)$. Since $\mathcal{Z}_Y$ is an affine curve with $|Y^{\mathfrak{T}}|$ components, and $p_2 : \mathcal{Z}_Y \rightarrow \mathbb{C}$ is a flat map whose restriction to each component is an isomorphism (by [4, Prop. 2]), the rank of $\mathbb{C}[\mathcal{Z}_Y]$ over $\mathbb{C}[v]$ is $|Y^{\mathfrak{T}}|$. Furthermore, by the long exact sequence of cohomology and the localization theorem, the rank of $H_{\mathfrak{T}}^{ev}(X, Y)$ is $|X^{\mathfrak{T}}| - |Y^{\mathfrak{T}}|$. But the rank of $\ker \overline{i_Y^*}$ is also $|X^{\mathfrak{T}}| - |Y^{\mathfrak{T}}|$. As shown above, $\ker \overline{i_Y^*} \subset \rho_X(H_{\mathfrak{T}}^{ev}(X, Y))$, so it follows that $\overline{i_Y^*}(\rho_X(H_{\mathfrak{T}}^{ev}(X, Y)))$ has rank zero in the free module $\mathbb{C}[\mathcal{Z}_Y]$, hence is trivial. Therefore, $I(\mathcal{Z}_Y) = \rho_X(H_{\mathfrak{T}}^{ev}(X, Y))$. To finish, we only need to show ψ_Y is injective. But this follows immediately from part (i). $\square$

The following example shows that non-principal subvarieties exist. Let

$$\lambda(t) = \begin{pmatrix} t^2 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & t^{-2} \end{pmatrix}$$

and

$$\varphi(v) = \begin{pmatrix} 1 & v & v^2/2 \\ 0 & 1 & v \\ 0 & 0 & 1 \end{pmatrix}.$$

Then $\lambda(t)\varphi(v)(\lambda(t))^{-1} = \varphi(t^2v)$, so λ and φ determine a two dimensional solvable subgroup of $SL_3(\mathbb{C})$, and hence a regular action $\mathfrak{B} \curvearrowright \mathbb{P}^2$. The $\mathfrak{T}$ fixed points are $o = [1, 0, 0]$, $\zeta_2 = [0, 1, 0]$, and $\zeta_3 = [0, 0, 1]$. Let w_1 and w_2 be the usual affine coordinates around o . Then the closures of $Y_1 = \{w_2 = 0\}$ and $Y_2 = \{2w_2 = w_1^2\}$ are the two $\mathfrak{B}$ -curves in $\mathbb{P}^2$. Let $Y = \overline{Y_1} \cup \overline{Y_2}$. Then Y is a $\mathfrak{B}$ -invariant subvariety such that $H^*(\mathbb{P}^2) \rightarrow H^*(Y)$ is not surjective. Indeed, $\dim H^2(\mathbb{P}^2) < \dim H^2(Y)$. On the other hand, $H_{\mathfrak{T}}^*(Y)$ is not generated by Chern classes of equivariant vector bundles. To see this, note that Y has vanishing odd cohomology, so one can use the localization theorem to compute its equivariant cohomology. In fact, by a well known result of Goresky, Kottwitz and MacPherson [9] (also see [3]), the image of $H_{\mathfrak{T}}^*(Y)$ in $H_{\mathfrak{T}}^*(Y^{\mathfrak{T}})$ consists of triples $(f_1(t), f_2(t), f_3(t))$ with all $f_i \in \mathbb{C}[t]$ such that $f_1(0) = f_2(0) = f_3(0)$. But not all classes of this form arise as polynomials in Chern classes of equivariant line bundles on Y , due to the fact that equivariance forces the further condition $f_2 = f_3 = -f_1$.

Remark. We do not know of an example of an irreducible $\mathfrak{B}$ -invariant subvariety of a $\mathfrak{B}$ -regular variety which is not principal.

6. EQUIVARIANT CHERN CLASSES

Let Y be a $\mathfrak{B}$ -invariant subvariety of a $\mathfrak{B}$ -regular variety X . The purpose of this section is to consider when the fundamental isomorphism $\rho_X : H_{\mathfrak{T}}^*(X) \rightarrow \mathbb{C}[\mathcal{Z}_X]$ defined in (4) can be defined for Y . For this, we need to consider $\mathfrak{B}$ -equivariant vector bundles on Y .

Recall that if V is an algebraic variety with an action of an algebraic group G , then a G -linearization of an algebraic vector bundle E on V is an action $G \times E \rightarrow E$ $((g, h) \rightarrow g \cdot h)$ such that for all $y \in V$, the restriction of $g \in G$ is a $\mathbb{C}$ -linear map $E_y \rightarrow E_{g \cdot y}$. In particular if $y \in V^G$, one obtains a representation of G on E_y and hence a representation of $\text{Lie}(G)$ on E_y . For $\xi \in \text{Lie}(G)$, we let ξ_y denote the corresponding endomorphism of E_y .

Suppose the vector bundle E admits a G -linearization. Recall that the k -th equivariant Chern class $c_k^G(E) \in H_G^{2k}(V)$ of E is the k -th Chern class of the vector bundle $E_G = (E \times \mathcal{E})/G$, where $\mathcal{E}$ is a contractible free G -space. The restriction of $c_k^G(E)$ to each $y \in V^G$ is the polynomial on $\text{Lie}(G)$ defined as

$$(7) \quad c_k^G(E)_y(\xi) = \text{Tr}_{\wedge^k E_y}(\xi_y).$$

We now turn to the case $G = \mathfrak{B}$, X $\mathfrak{B}$ -regular and Y a $\mathfrak{B}$ -invariant subvariety. Put

$$\mathcal{W} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \mathcal{V} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}.$$

By [4, Lemma 1], if E is $\mathfrak{B}$ -equivariant and $(y, v) \in \mathcal{Z}_X$, then

$$\rho_X(c_k^{\mathfrak{B}}(E))(y, v) = \text{Tr}_{\wedge^k E_y}(v\mathcal{W} - 2\mathcal{V})_y,$$

where $(v\mathcal{W} - 2\mathcal{V})_y$ is the endomorphism determined by $v\mathcal{W} - 2\mathcal{V} \in \text{Lie}(\mathfrak{B})$. This is a key fact since the smoothness of X implies $H_{\mathfrak{T}}^*(X)$ is generated by Chern classes of $\mathfrak{B}$ -equivariant vector bundles on X [5, Prop. 3].

Let $\mathcal{H}_{\mathfrak{B}}^*(Y)$ denote the subalgebra of $H_{\mathfrak{T}}^*(Y)$ generated by equivariant Chern classes of $\mathfrak{B}$ -equivariant vector bundles on Y . As just noted, $\mathcal{H}_{\mathfrak{B}}^*(X) = H_{\mathfrak{T}}^*(X)$. The goal of the remainder of this section is to prove that if E is a $\mathfrak{B}$ -linearized vector bundle on Y then $\rho_Y(c_k^{\mathfrak{B}}(E))$ is a regular function on the affine curve $\mathcal{Z}_Y$. Thus, there exists a well defined map

$$\rho_Y : \mathcal{H}_{\mathfrak{B}}^*(Y) \rightarrow \mathbb{C}[\mathcal{Z}_Y]$$

such that the diagram

$$(8) \quad \begin{array}{ccc} H_{\mathfrak{T}}^*(X) & \xrightarrow{\rho_X} & \mathbb{C}[\mathcal{Z}_X] \\ i_Y^* \downarrow & & \downarrow \overline{i_Y^*} \\ \mathcal{H}_{\mathfrak{B}}^*(Y) & \xrightarrow{\rho_Y} & \mathbb{C}[\mathcal{Z}_Y]. \end{array}$$

is commutative.

Before beginning the proof, assume $v \neq 0$ and define

$$\mathfrak{H}_v = \varphi(1/v)\mathfrak{T}\varphi(-1/v).$$

By (3), $(y, v) \in \mathcal{Z}_Y$ if and only if $y \in Y^{\mathfrak{H}_v}$. Also, put

$$\mathcal{H} = \overline{\bigcup_{v \neq 0} \mathfrak{H}_v \times \{v\}} \subset \mathfrak{B} \times \mathbb{C}.$$

Lemma 9. *Let $\pi : \mathcal{H} \rightarrow \mathbb{C}$ be projection on the second factor. Then:*

$$\pi^{-1}(0) = \mathfrak{U} \cup -(\mathfrak{U}).$$

Proof. First note the identity

$$(9) \quad \begin{pmatrix} 1 & v^{-1} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \begin{pmatrix} 1 & -v^{-1} \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} a & (1 - a^2)(av)^{-1} \\ 0 & a^{-1} \end{pmatrix}.$$

One must find all the possible finite limits of the right-hand side as $v \rightarrow 0$. Take a sequence $(a_i, v_i) \neq (0, 0)$ such that $v_i \rightarrow 0$, and the right-hand side of (9) has a finite limit. In order for this to happen, $(1 - a_i^2)/a_i \rightarrow 0$, so $a_i^2 \rightarrow 1$. Hence, if a limit exists, it lies in $\mathfrak{U} \cup -(\mathfrak{U})$. The reverse inclusion is similar. $\square$

Consequently, if we put $\mathfrak{h}_v = \text{Lie}(\mathfrak{H}_v)$ for $v \neq 0$ and set $\mathfrak{h}_0 = \text{Lie}(\mathfrak{U})$, then the family of Lie algebras of the fibres of $\mathcal{H}$ is

$$\mathfrak{h} = \bigcup_{v \in \mathbb{C}} \mathfrak{h}_v \times \{v\} \subset \text{Lie}(\mathfrak{B}) \times \mathbb{C},$$

and, moreover, $v\mathcal{W} - 2\mathcal{V}$ is a non-vanishing regular section of π . For convenience, let $s(v) = v\mathcal{W} - 2\mathcal{V}$.

Now let E be a $\mathfrak{B}$ -linearized vector bundle on Y , and let $\tilde{E}$ be the pull-back of E to $Y \times \mathbb{C}$. Assume $v \in \mathbb{C}$ and $y \in Y^{\mathfrak{H}_v}$. That is, $(y, v) \in \mathcal{Z}_Y$. Then E_y admits an $\mathfrak{H}_v$ -representation and hence an $\mathfrak{h}_v$ -representation. For $\xi \in \mathfrak{h}_v$ let ξ_y denote the corresponding endomorphism on E_y . Then the upshot of our discussion is the following:

Lemma 10. *The map from $\mathcal{Z}_Y$ to $\text{End}(\tilde{E}|_{\mathcal{Z}_Y})$ given by*

$$(y, v) \mapsto s(v)_y,$$

is a regular section of the bundle $\text{End}(\tilde{E}|_{\mathcal{Z}_Y})$. Consequently the map

$$(y, v) \mapsto \text{Tr}_{\bigwedge^k E_y}(s(v)_y)$$

is a regular function on $\mathcal{Z}_Y$.

Lemma 11. *Let E be a $\mathfrak{B}$ -linearized vector bundle on Y . Then $\rho_Y(c_k^{\mathfrak{T}}(E))$ is a regular function on $\mathbb{C}[\mathcal{Z}_Y]$.*

Proof. View v as an element of $\text{Lie}(\mathfrak{T})$. As above let v_{ζ_j} denote the corresponding endomorphism on E_{ζ_j} . By Equation (7) we have

$$c_k^{\mathfrak{T}}(E)_{\zeta_j}(v) = \text{Tr}_{\bigwedge^k E_y}(v_{\zeta_j}).$$

Put $\zeta_j(v) = \varphi(v^{-1})\zeta_j$. Then the $\zeta_j(v)$ are the fixed points of $\mathfrak{H}_v$. Recall that $s(v) \in \mathfrak{h}_v$, hence it gives an endomorphism $s(v)_{\zeta_j(v)}$ of the fibre $E_{\zeta_j(v)}$. Since E is $\mathfrak{B}$ -linearized, the element $\varphi(v^{-1})$ gives an isomorphism $E_{\zeta_j} \rightarrow E_{\zeta_j(v)}$; moreover the endomorphism $v_{\zeta_j(v)}$ on $E_{\zeta_j(v)}$ is conjugate to the endomorphism v_{ζ_j} on E_{ζ_j} . Put $y = \zeta_j(v)$. Then $(y, v) \in \mathcal{Z}_Y$ and

$$\text{Tr}_{\bigwedge^k E_{\zeta_j}}(v_{\zeta_j}) = \text{Tr}_{\bigwedge^k E_y}(s(v)_y).$$

But by Lemma 10, the function $(y, v) \mapsto \text{Tr}_{\bigwedge^k E_y}(s(v)_y)$ is a regular function on $\mathcal{Z}_Y$. $\square$

To summarize the discussion of this section, we state

Theorem 12. *If Y is a $\mathfrak{B}$ -invariant subvariety of a $\mathfrak{B}$ -regular variety X , then we obtain a $\mathbb{C}[v]$ -algebra homomorphism $\rho_Y : \mathcal{H}_{\mathfrak{B}}^*(Y) \rightarrow \mathbb{C}[\mathcal{Z}_Y]$ such that the diagram (8) is commutative.*

7. CLASSIFICATION OF PRINCIPAL SUBVARIETIES

We now classify principal subvarieties. First, we describe the image of $H_{\mathfrak{T}}^*(X)$ in $H_{\mathfrak{T}}^*(Y)$ for a $\mathfrak{B}$ -invariant subvariety Y with vanishing odd cohomology.

Theorem 13. *Suppose Y is $\mathfrak{B}$ -invariant subvariety of a $\mathfrak{B}$ -regular variety X with vanishing odd cohomology. Then*

(i) $\rho_Y : \mathcal{H}_{\mathfrak{B}}^*(Y) \rightarrow \mathbb{C}[\mathcal{Z}_Y]$ is an isomorphism, and

(ii) $i_Y^*(H_{\mathfrak{T}}^*(X)) = \mathcal{H}_{\mathfrak{B}}^*(Y)$.

Consequently, $\mathcal{H}_{\mathfrak{B}}^*(Y)$ is exactly the subalgebra generated by Chern classes of $\mathfrak{B}$ -equivariant vector bundles on Y which are pull backs of $\mathfrak{B}$ -equivariant vector bundles on X .

Proof. It is clear from the commutativity of (8) that ρ_Y is surjective. The definition of ρ_Y and localization (applied to Y) implies it is injective. This proves (i). We now show (ii). By Lemma 8, we have an isomorphism $\psi_Y : \mathbb{C}[Z_Y] \rightarrow i_Y^*(H_{\mathfrak{T}}^*(X))$. We claim that $\rho_Y \psi_Y = 1$. In fact, by (8) and the definition of ψ_Y ,

$$\rho_Y \psi_Y = \rho_Y i_Y^* \rho_X^{-1} \overline{i_Y^*}^{-1} = \overline{i_Y^*} \rho_X \rho_X^{-1} \overline{i_Y^*}^{-1},$$

which is clearly the identity. Thus

$$\rho_Y(i_Y^*(H_{\mathfrak{T}}^*(X))) = \mathbb{C}[Z_Y],$$

which certainly implies (ii). □

Corollary 14. *A $\mathfrak{B}$ -invariant subvariety Y of a $\mathfrak{B}$ -regular variety X is principal if and only if $\mathcal{H}_{\mathfrak{B}}^*(Y) = H_{\mathfrak{T}}^*(Y)$.*

Proof. The necessity is clear. But if $\mathcal{H}_{\mathfrak{B}}^*(Y) = H_{\mathfrak{T}}^*(Y)$, then Y has vanishing odd cohomology, so the result follows from Theorem 13 (b). □

Theorem 2 follows immediately from this corollary. We conclude with an application.

Corollary 15. *Suppose Y is a normal $\mathfrak{B}$ -invariant subvariety of a $\mathfrak{B}$ -regular variety X such that $H^*(Y)$ is generated by Chern classes of line bundles. Then Y is principal.*

Proof. Let L be an algebraic line bundle on Y . By [10], some power L^m is $\mathfrak{B}$ -equivariant. Hence the first Chern classes of $\mathfrak{B}$ -equivariant line bundles on Y generate $H^*(Y)$. This is equivalent to saying that $\mathcal{H}_{\mathfrak{B}}^*(Y) = H_{\mathfrak{T}}^*(Y)$. □

Acknowledgement: We would like to thank Michel Brion and Jochen Kuttler who each noticed an error in a previous version of this paper.

REFERENCES

- [1] E. Akyildiz and J.B. Carrell: *Cohomology of projective varieties with regular SL_2 actions*. Manuscripta Math. **58** (1987), 473–486.
- [2] E. Akyildiz and J.B. Carrell: *A generalization of the Kostant–Macdonald identity*, Proc. Nat. Acad. Sci. U.S.A. **86** (1989), 3934–3937.
- [3] M. Brion: *Equivariant cohomology and equivariant intersection theory*. Notes by Alvaro Rittatore. NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., 514, Representation theories and algebraic geometry (Montreal, PQ, 1997), 1–37, Kluwer Acad. Publ., Dordrecht, 1998.
- [4] M. Brion and J.B. Carrell: *The equivariant cohomology ring of regular varieties*. Michigan Math. J. **52** (2004), 189–203.
- [5] J. B. Carrell: *Deformation of the nilpotent zero scheme and the intersection rings of invariant subvarieties*, J. Reine Angew. Math. **460** (1995), 37–54.
- [6] J. B. Carrell and D. I. Lieberman: *Holomorphic vector fields and compact Kaehler manifolds*. Invent. Math. **21** (1973), 303–309.
- [7] J. B. Carrell and D. I. Lieberman: *Vector fields and Chern numbers*. Math. Ann. **225** (1977), 263–273.
- [8] C. De Concini and C. Procesi: *Complete symmetric varieties*. Invariant theory (Montecatini, 1982), 1–44, Lecture Notes in Math., **996**, Springer, Berlin, 1983.
- [9] M. Goresky, R. Kottwitz and R. MacPherson: *Equivariant cohomology, Koszul duality, and the localization theorem*. Invent. Math. **131** (1998), 25–83.
- [10] F. Knop, H. Kraft, D. Luna and T. Vust: *Local properties of algebraic group actions*. Algebraische Transformationsgruppen und Invariantentheorie, 63–75, DMV Sem. **13**, Birkhäuser, Basel, 1989.
- [11] N. Spaltenstein: *The fixed point set of a unipotent transformation on the flag manifold*. Indag. Math. **38** (1976), 452–456.
- [12] J. Tymoczko: *Paving Hessenberg varieties by affines*. Selecta Math. (N.S.) **13** (2007), 353–367.

James B. Carrell, University of British Columbia, Vancouver, B.C., Canada
Email address: carrell@math.ubc.ca

Kiumars Kaveh, University of Toronto, Toronto, ON., Canada
Email address: kaveh@math.toronto.edu